

A Bi-Photon Quantum Interferometer for Complete Measurement of the Bi-Photon Wave Function

Roey Pomerantz, Yaakov Shaked and Avi Pe'er¹

¹*Department of physics and BINA Center of nano-technology, Bar-Ilan university, Ramat-Gan 52900, Israel**

We demonstrate a simple two-photon, nonclassical interference between the generation amplitudes of ultra-broadband entangled photon pairs (bi-photons) in two different nonlinear media, allowing measurement of the bi-photons spectral phase. In an analogy to a Mach-Zehnder interferometer, the nonlinear media act as beam splitters that couple between the pump beam and the bi-photons. The nonclassical nature of the interference is demonstrated by observing the dependence of the interference contrast on internal loss, analogous to an attempt to obtain "which-path" information.

PACS numbers: 42.50.Dv, 42.65.Lm, 42.40.Kw

Due to quantum correlation, the state of a bi-photon (entangled photon pair) is defined well beyond the uncertainty regarding each of the constituent photons. The inherent quantum nature of bi-photons is a foundation in quantum optics, exploited for many experiments and applications, such as verification of quantum theory [1–4], engineering of Bell states for quantum information [5–9] and sources of squeezed light for measurements of optical phase below the shot-noise limit [10–12]. A most pronounced realizations of this quantum correlation is with ultra-broadband time-energy entangled bi-photons, produced by type-I spontaneous parametric down conversion (SPDC) from a narrowband pump laser. The precise energy-sum correlation of broadband bi-photons can extend over nearly an octave in the spectrum (more than 100THz in this report), and their time-difference correlation can be in the few fs regime [13–17]. As a result of this ultrashort time correlation, a very high flux of bi-photons (up to 10^{14} bi-photons per second) can be generated and still be composed of single bi-photons, with negligible probability for multiple pairs [15, 17, 18].

In spite of their unique quantum properties, broadband bi-photons are rarely used in quantum information applications, mainly because of the bandwidth incompetency in the standard detection scheme between the bi-photons and the photo-detectors. When the single photons are directly detected, the photo-detectors response time is far too slow (~ 100 ps with the fastest available detectors) to resolve the ultrafast correlation time of the bi-photons (of order $10 - 100$ fs). In frequency domain, this incompetency leads to an undesired distinguishability between different frequency pairs of the bi-photons spectrum. As a result, the broadband spectrum of bi-photons is a burden, not a resource with standard detection, and much effort is invested in many experiments to eliminate time energy entanglement by matching the bi-photons bandwidth to that of the pump, either by narrowing of the bi-photons [19–21] or by increasing the pump bandwidth using ultrashort pump pulses [22–24].

In order to exploit the bandwidth resource of bi-photons, a different route for detection is required, where

the frequency pairs of the bi-photons remain undistinguished. In the past, broadband physical two-photon detection, such as sum-frequency generation (SFG) was used for this purpose [17, 18, 25, 26]. In SFG, all frequency pairs that sum up to the original pump interfere in the SFG generation, leading to the desired indistinguishability. The quantum nature of the interference with bi-photons was demonstrated by a linear dependence of the nonlinear SFG intensity on the incoming bi-photons flux [18].

SFG indeed offers true broadband two-photon detection, but with two major limitations: first, the low efficiency of the SFG process in the single photon level ($\sim 10^{-10} - 10^{-8}$) yields a very low flux of SFG photons that is difficult to detect, especially against the background from scattered pump light. Second, as a broadband pairwise interference effect, the SFG signal is highly sensitive to spectral phase modulations of the bi-photon input. SFG can therefore detect only nearly transform limited bi-photons and exact dispersion compensation is required to obtain a detectable SFG signal. By homodyne detection of the generated SFG with respect to the original pump, the overall bi-photon phase can be measured [25, 27], but not the spectral phase of the composing frequency-pairs.

Here we demonstrate a simple interferometric measurement of the spectral phase (and amplitude) of bi-photons. The experimental concept, schematically outlined in Fig. 1(a), includes two nonlinear crystals in series that are pumped by the same pump at frequency Ω_p , where the spectrum of bi-photons is measured after the second crystal. Bi-photons produced in the first crystal enter the second crystal, where due to the presence of the pump they can either stimulate further down conversion, or be up-converted back to the pump, depending on the relative phase between the pump and the bi-photons [28–30]. If the bi-photons are not transform limited, the relative pump - bi-photons phase varies spectrally, leading to the appearance of high-contrast interference fringes on the measured bi-photons spectrum in a symmetric manner around the degeneracy point at $\Omega_p/2$.




FIG. 1: (color online) (a) A simplified block diagram of the experimental process, showing the generation of bi-photons by SPDC in the first crystal, a change of the relative phase between the pump and the down converted light after the crystal and another generation of down converted light in the 2nd crystal. (b) The two-photon Mach-Zehnder analog of the experiment: Each crystal is analogous to a beam splitter (unbalanced) that couples between the bi-photons and pump beams, allowing measurement of the relative spectral phase between the bi-photons and pump.

This spectral interference provides a direct holographic measurement of the bi-photons spectral phase. Since bi-photons can be generated either in the first crystal or the second, quantum interference between the probability amplitudes of those two indistinguishable possibilities occurs according to the relative phase between the pump and the bi-photons. The described setup is therefore analogous to a Mach-Zehnder interferometer for the bi-photons, where the crystals represent beam splitters that couple the pump and the bi-photons beams as illustrated in fig. 1 (b).

While the mere existence of the interference can be explained by classical theory for high-gain parametric amplification, the observed fringe contrast with bi-photons is a clear quantum signature [28–30]. We demonstrate the nonclassical nature by comparing two experimental situations that are classically indistinguishable, yet quantum mechanically lead to very different results: we attenuate the light fields entering the 2nd crystal in two ways - once by attenuating the pump before the first crystal, thereby attenuating also the flux of bi-photons generated in it, and second by attenuating both the pump and the bi-photons beam between the two crystals. Quantum-mechanically these two possibilities are different procedures - the first attenuates linearly the generation rate of bi-photons, but does not alter their cor-

relation (two-photon attenuation), preserving the fringes contrast, whereas the second attenuates every photon independently (one-photon attenuation), thereby reducing *quadratically* the flux of "surviving" bi-photons that enter the second crystal, leading to a linear reduction of the fringe contrast in the bi-photon interference. In the Mach-Zehnder analog, attenuation between the crystals is equivalent to an attempt to obtain "which path" information, causing the interference to disappear [31].




FIG. 2: (a) Experimental layout: bi-photons are generated in the 1st crystal (12mm long PPKTP), pumped by a single-frequency diode laser at 880 nm with up to 400 mW power. Both the bi-photons and the pump are directed into a second identical crystal, where further generation of bi-photons or up-conversion back to the pump can occur. The resulting bi-photons spectrum is measured by a home-built spectrometer composed of a prism (SF11) and a CCD camera (Xeva-2.5-320 by Xenics). The last mirror M4 separates the pump from the bi-photons, allowing the pump power to be measured. Attenuation is achieved by a half-wave plate and polarizer, either before the 1st crystal or between the crystals. (b) Calculated phase mismatch for collinear down-conversion in the PPKTP crystals, indicating ultra-broad phase matching around the zero-dispersion of KTP. (c) SPDC spectrum from a single crystal, demonstrating bi-photons spanning nearly an octave.

A simple quantum model accounts for the observed bi-photon interference. The bi-photon state $|\Psi\rangle$ created by the first crystal can be approximated as:

$$|\Psi\rangle_{1DC} \approx |0\rangle + \int_0^\infty d\omega g(\omega) |1_{\omega_0+\omega}, 1_{\omega_0-\omega}\rangle \quad (1)$$

where $|0\rangle$ is the vacuum input of the 1st crystal, ω_0 is the frequency of the center of the spectrum and $|g(\omega)|^2$ represents the spectral photon number density determined by the pump intensity and the phase-matching conditions

in the down conversion crystal. Assuming $|g(\omega)|^2 \ll 1 \text{ Hz}^{-1}$, higher order terms of multiple bi-photons can be neglected (in our experiment $|g(\omega)|^2 \sim 1/90 \text{ Hz}^{-1}$). In the Mach-Zehnder analog, $|g(\omega)|$ represents the reflection/transmission of the BS.

The two photon state entering the second crystal after experiencing spectral phase modulation $\phi(\omega)$ (e.g. from material / mirror dispersion) is

$$|\Psi\rangle_{1DC} \approx |0\rangle + \int_0^\infty d\omega g(\omega) e^{i\Phi(\omega)} |1_{\omega_0+\omega}, 1_{\omega_0-\omega}\rangle \quad (2)$$

where $\Phi(\omega) = \phi(\omega_0 - \omega) + \phi(\omega_0 + \omega)$ is the spectral phase of the bi-photon - the sum of phases of twin frequency pairs. Neglecting multiple pairs generation also in the second crystal, the quantum state after the second crystal is

$$|\Psi\rangle_f \approx |0\rangle + \int_0^\infty d\omega g(\omega) \left(1 + e^{i\Phi(\omega)}\right) |1_{\omega_0+\omega}, 1_{\omega_0-\omega}\rangle \quad (3)$$

and the detected spectral intensity of bi-photons equals

$$I(\omega) = \langle \Psi_f | a^\dagger a | \Psi_f \rangle \propto |g(\omega)|^2 \cos^2(\Phi(\omega)/2). \quad (4)$$

A detected down-converted photon at frequency ω can be generated either in the 1st crystal or in the 2nd, and the interference between those two probability amplitudes leads to the observed fringes. Constructive interference represents enhanced down-conversion in the second crystal, whereas destructive interference represents up-conversion of bi-photons back to the pump.

In the experiment, described in Fig.2 (a), a single frequency diode laser at 880nm pumps a KTP crystal periodically polled for collinear down conversion around 1760nm . This pump wavelength was chosen to match the center of the bi-photons spectrum to the zero-dispersion wavelength of the KTP crystal, allowing an ultra-broad bandwidth (over 100 THz , nearly an octave) for collinear down conversion between $1.3 - 2.5 \mu\text{m}$, as illustrated in Fig. 2(b). Figure 2(c) shows the measured bi-photons spectrum. Such bandwidth corresponds to a maximum possible bi-photon flux of $F_{max} = 1.08 \times 10^{14}$ photons/s, nearly $12 \mu\text{W}$ of single photon pairs. The actual flux in the experiment was approximately $F_{max}/90$, limited by the available pump power, well within the single bi-photon regime. The down converted light from the first KTP crystal continues along with the pump into a second identical KTP crystal, where either down conversion or up conversion back to the pump can occur. Negative dispersion of the broadband dielectric mirrors (M2, M3) between the crystals introduces a spectral modulation of the phase between the different frequency pairs of the down conversion and the pump, causing interference fringes to appear on the bi-photons spectrum, as shown in Fig. 3. An attenuator is inserted either before the first crystal or after it, and the down conversion spectrum is measured by a prism-based spectrometer.




FIG. 3: (color online) Spectral fringes (a) Raw CCD image of spectral fringes after both crystals. (b) Calibrated spectrum of bi-photons light after both crystals (red line, marked 2), with only one crystal (dashed blue line, marked 1), and the corresponding calculated spectral phase $\Phi(\omega)/2$ of the bi-photons (dotted green line marked 3).

The observed fringe contrast in our experiment of $\sim 20\%$ (calculated by averaging the contrast of the central three fringes) is limited both technically and fundamentally. Technical limits are due to residual non-collinear generation from high order spatial components of the pump beam. The fundamental limit is due to the use of focused Gaussian beams in the finite-length crystals. Due to the focusing into the crystals (necessary for high bi-photon generation), both the pump and the bi-photons incur a Gouy phase shift [32] across each focus. Since in collinear down-conversion the Rayleigh range of the pump and the bi-photons is equal, the incur-rate of the Gouy phase for bi-photons is double that of the pump. The relative pump to bi-photons phase therefore varies across the focus within the second crystal, thus reducing the fringe contrast according to the ratio of crystal length to the Rayleigh range.

Figure 4 presents the measured fringe contrast as function of pump (and bi-photons) power entering the second crystal for the two cases of attenuation. First, before the first crystal (two-photon attenuation), which reduces the bi-photon flux, but does not affect the correlation, and therefore leaves the fringe contrast intact. Second, between the crystals (one-photon attenuation), which reduces linearly the quantum correlation between the photon pairs, causing a linear reduction of the observed contrast. This clear difference between two classically indistinguishable scenarios is a signature of the quantum nature of the interference.




FIG. 4: (color online) Measured fringe contrast as function of attenuation. (a) Fringe contrast with two-photon attenuation before the first crystal (blue points) and with one-photon attenuation between the crystals (red points). Sample spectral interferogram are shown in (b) and (c) for two-photon and one-photon attenuation correspondingly at 100mW pump level (25% transmission), demonstrating the reduction in fringe contrast with one-photon attenuation.

We note that the dark fringes of the spectral interferogram represent up-conversion of single bi-photons back to the pump with 20% efficiency (!), orders of magnitude higher than with direct SFG [17, 18]. This phenomenal enhancement is due to the large gain offered by the strong pump in the second crystal, a two-photon analog of the homodyne gain from a strong local oscillator in homodyne detection.

In conclusion, we demonstrated a simple, general and robust method for holographic measurement of the spectral phase (and amplitude) of entangled broadband bi-photons through non-classical two-photon interference, and to measure the purity of the two-photon state through the interference fringe contrast. The ultra-high flux of bi-photons and the high two-photon interaction efficiency of single bi-photons, enhanced orders of magnitude (ideally to unity) due to the intense pump field, simplify the measurement compared to standard experiments at the single photon level, allowing measurement of the bi-photons with a simple camera. Due to the collinear arrangement, the experimental configuration is inherently robust to phase fluctuations and does not require any phase locking to observe a stable fringe pattern. We expect this method to become an important member of the quantum optics tool box for broadband time-energy entangled photons.

* Electronic address: avi.peer@biu.ac.il

- [1] A. Aspect, P. Grangier, and G. Roger, *Phys. Rev. Lett.* **49**, 91 (1982).
- [2] Z. Y. Ou, S. F. Pereira, and H. J. Kimble, *Appl. Phys. B* **55**, 265 (1992).
- [3] P. G. Kwiat, K. Mattle, H. Weinfurter, A. Zeilinger, A. V. Sergienko, and Y. Shih, *Phys. Rev. Lett.* **75**, 4337 (1995).
- [4] J. C. Howell, R. S. Bennink, S. J. Bentley, and R. W. Boyd, *Phys. Rev. Lett.* **92**, 210403 (2004).
- [5] D. M. Greenberger, M. A. Horne, A. Shimony, and A. Zeilinger, *American Journal of Physics* **58**, 1131 (1990).
- [6] H. J. Briegel and R. Raussendorf, *Phys. Rev. Lett.* **86**, 910 (2001).
- [7] E. Megidish, T. Shacham, A. Halevy, L. Dovrat, and H. S. Eisenberg, *Phys. Rev. Lett.* **109**, 080504 (2012).
- [8] N. C. Menicucci, S. T. Flammia, and O. Pfister, *Phys. Rev. Lett.* **101**, 130501 (2008).
- [9] M. Pysher, Y. Miwa, R. Shahrokhshahi, R. Bloomer, and O. Pfister, *Phys. Rev. Lett.* **107**, 030505 (2011).
- [10] C. M. Caves, *Phys. Rev. D* **23**, 1693 (1981).
- [11] M. J. Holland and K. Burnett, *Phys. Rev. Lett.* **71**, 1355 (1993).
- [12] T. Kim, O. Pfister, M. J. Holland, J. Noh, and J. L. Hall, *Phys. Rev. A* **57**, 4004 (1998).
- [13] C. K. Hong, Z. Y. Ou, and L. Mandel, *Phys. Rev. Lett.* **59**, 2044 (1987).
- [14] I. Abram, R. K. Raj, J. L. Oudar, and G. Dolique, *Phys. Rev. Lett.* **57**, 2516 (1986).
- [15] S. E. Harris, *Phys. Rev. Lett.* **98**, 063602 (2007).
- [16] B. Dayan, A. Pe'er, A. A. Friesem, and Y. Silberberg, *Phys. Rev. Lett.* **93**, 023005 (2004).
- [17] A. Pe'er, B. Dayan, A. A. Friesem, and Y. Silberberg, *Phys. Rev. Lett.* **94**, 073601 (2005).
- [18] B. Dayan, A. Pe'er, A. A. Friesem, and Y. Silberberg, *Phys. Rev. Lett.* **94**, 043602 (2005).
- [19] A. Christ and C. Silberhorn, *Phys. Rev. A* **85**, 023829 (2012).
- [20] A. Christ, A. Eckstein, P. J. Mosley, and C. Silberhorn, *Optics Express* **17**, 3441 (2009).
- [21] Y.-P. Huang, J. B. Altepeter, and P. Kumar, *Phys. Rev. A* **84**, 033844 (2011).
- [22] W. P. Grice and I. A. Walmsley, *Phys. Rev. A* **56**, 16271634 (1997).
- [23] P. J. Mosley, J. S. Lundeen, B. J. Smith, P. Wasylczyk, A. B. U'Ren, C. Silberhorn, and I. A. Walmsley, *Phys. Rev. Lett.* **100**, 133601 (2008).
- [24] J. F. Hodelin, G. Khoury, and D. Bouwmeester, *Phys. Rev. A* **74**, 013802 (2006).
- [25] S. Sensarn, I. Ali-Khan, G. Y. Yin, and S. E. Harris, *Phys. Rev. Lett.* **102**, 053602 (2009).
- [26] N. P. Georgiades, E. S. Polzik, K. Edamatsu, H. J. Kimble, and A. S. Parkins, *Phys. Rev. Lett.* **75**, 3426 (1995).
- [27] C. Belthangady, S. Du, C.-S. Chu, G. Y. Yin, and S. E. Harris, *Phys. Rev. A* **80**, 031803 (2009).
- [28] T. J. Herzog, J. G. Rarity, H. Weinfurter, and A. Zeilinger, *Phys. Rev. Lett.* **72**, 629 (1994).
- [29] X. Y. Zou, L. J. Wang, and L. Mandel, *Phys. Rev. Lett.* **67**, 318 (1991).
- [30] I. R. Senitzky, *Phys. Rev. Lett.* **73**, 3040 (1994).
- [31] B.-G. Englert, *Phys. Rev. Lett.* **77**, 2154 (1996).

- [32] A. E. Siegman, *Lasers* (University Science Books, 1986).